

11-286-C025

Monoclonal Antibody to 5-bromodeoxyuridine (BrdU) Purified Antibody (0.025 mg)

Clone:	MoBu-1
Isotype:	Mouse IgG1
Specificity:	The antibody MoBu-1 reacts specifically with BrdU incorporated into DNA during S-phase of a cell cycle. The antibody MoBu-1 is also useful for detecting proliferating cells by flow cytometry or immunofluorescence staining. It reacts also specifically with 5-bromouridine (BrU).
Regulatory Status:	RUO
Immunogen:	5-bromodeoxyuridine conjugated with hemocyanine.
Application:	Immunohistochemistry (paraffin sections) Application note: excellent Flow Cytometry Recommended dilution: 1-2 µg/ml Immunocytochemistry Recommended dilution: 2 µg/ml
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified from ascites by protein-A affinity chromatography.
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	Bromodeoxyuridine (BrdU) is a thymidine analog which is selectively incorporated into the DNA of proliferating cells to provide a marker for the DNA being replicated. The number of proliferating cells can then be detected in cell lysates, tissue sections or suspensions using an antibody specific for the BrdU.
References:	<p>*Ashby J, Tinwell H, Soames A, Foster J: Induction of hyperplasia and increased DNA content in the uterus of immature rats exposed to coumestrol. Environ Health Perspect. 1999 Oct;107(10):819-22.</p> <p>*Soames AR, Lavender D, Foster JR, Williams SM, Wheeldon EB: Image analysis of bromodeoxyuridine (BrdU) staining for measurement of S-phase in rat and mouse liver. J Histochem Cytochem. 1994 Jul;42(7):939-44.</p> <p>*Buckiova D, Kubinova L, Soukup A, Jelinek R, Brown NA: Hyperthermia in the chick embryo: HSP and possible mechanisms of developmental defects. Int J Dev Biol. 1998 Jul;42(5):737-40.</p> <p>*Stanek D, Kiss T, Raska I: Pre-ribosomal RNA is processed in permeabilised cells at the site of transcription. Eur J Cell Biol. 2000 Mar;79(3):202-7.</p>

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 42 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz